

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 132 086 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
12.09.2001 Bulletin 2001/37

(51) Int Cl.⁷: **A61K 31/4015, A61P 13/12**

(21) Application number: **01300690.3**

(22) Date of filing: **25.01.2001**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
AL LT LV MK RO SI

• **Thomson, David Duane, Pfizer Global
Groton, Connecticut 06340 (US)**

(30) Priority: **31.01.2000 US 178968 P**

(71) Applicant: **Pfizer Products Inc.
Groton, Connecticut 06340 (US)**

(74) Representative: **McMunn, Watson Palmer et al
Pfizer Limited
Patents Department
Ramsgate Road
Sandwich, Kent CT13 9NJ (GB)**

(72) Inventors:
• **Paralkar, Vishwas Madhav, Pfizer Global
Groton, Connecticut 06340 (US)**

Remarks:

A request for a correction in claim 03 has been filed pursuant to Rule 88 EPC. A decision on the request will be taken during the proceedings before the Examining Division (Guidelines for Examination in the EPO, A-V, 3.).

(54) **Use of prostaglandin (PGE₂) receptor 4 (EP₄) selective agonists for the treatment of acute and chronic renal failure**

(57) This invention is directed to methods and compositions of treating acute or chronic renal failure or dys-

function, or conditions caused thereby comprising administering prostaglandin agonists, which are EP₄ receptor selective prostaglandin agonists.

EP 1 132 086 A2

Description**FIELD OF THE INVENTION**

5 **[0001]** The present invention relates to methods and pharmaceutical compositions comprising receptor selective prostaglandin (PGE₂) agonists for the treatment of kidney diseases, such as chronic and acute renal failure or dysfunction, in animals, particularly mammals. More specifically, the present invention relates to such methods and pharmaceutical compositions comprising type 4 (EP₄) receptor selective prostaglandin (PGE₂) agonists.

10 BACKGROUND OF THE INVENTION

[0002] The naturally occurring prostaglandins are comprised of several biological entities including PBD, PGE, PGF, PGG, PGH and PGI. It has been well documented that prostaglandins have effects on many of the organs and systems of the body.

15 **[0003]** In the kidney, the prostaglandins modulate renal blood flow and may serve to regulate urine formation by both renovascular and tubular effects. In clinical studies, PGE₁ has been used to improve creatinine clearance in patients with chronic renal disease, to prevent graft rejection and cyclosporine toxicity in renal transplant patients, to reduce the urinary albumin excretion rate and N-acetyl-beta-D-glucosaminidase levels in patients with diabetic nephropathy, and to improve urea clearance in healthy volunteers. PGE₁ also has been administered intravenously during surgeries

20 **[0004]** Renal dysfunction and/or renal failure is manifested in the body in a number of different ways. Any one or a combination of the following manifestations could indicate renal dysfunction or failure in a patient: lower than normal creatinine clearance; lower than normal free water clearance; higher than normal blood urea and/or nitrogen and/or potassium and/or creatinine levels; altered activity of kidney enzymes such as gamma glutamyl synthetase, alanine phosphatidase, N-acetyl-beta-D-glucosaminidase, or beta-2-microglobulin; altered urine osmolarity or volume; increase over normal levels or new observation of microalbuminuria or macroalbuminuria; or need for dialysis. Successful prevention of renal dysfunction or renal failure is indicated if the above described events do not occur at all, if they occur with less severity, if they occur in fewer patients at risk for renal dysfunction or renal failure; or if the patient recovers from these problems more quickly than normal.

30 **[0005]** Acute renal failure caused by the injection of contrast media has been recognized for many years as a complication of procedures utilizing such media. It has been estimated that the incidence of acute renal failure directly induced by contrast media is 10-15%, while the incidence of contrast associated nephropathy defined by clinically significant increases in serum creatinine is as high as 22%. See Porter, Am. J. Cardiol., 64: 22E-26E (1989). U.S. Patent No. 5,807,895 discloses a method of preventing renal failure or dysfunction caused by medical procedures which utilize contrast media by intravenous administration of a prostaglandin compound selected from PGE₁, PGE₂, PGI₂ or an analog or pharmaceutically acceptable salt thereof.

40 **[0006]** Chronic renal failure (CRF) occurs as a result of progressive and later, permanent reduction in the glomerular filtration rate (GFR), which is associated with loss of functional nephron units. When the GFR continues to decline to less than 10% of normal (5-10 ml/min), the subject progresses to end-stage renal failure (ESRD). R.A. Lafayette, R. D. Perrone and A.S. Levey: "Laboratory Evaluation of Renal Function," in: Diseases of the Kidney (Eds: R.W. Schrier and C.W. Gottschalk), Little, Brown and Company, Inc., Vol. 6, 307-354 (1997). At this point, unless the subject receives renal replacement therapy (i.e., chronic hemodialysis, continued peritoneal dialysis or kidney transplantation) renal failure will rapidly progress to cause death. It is believed that the therapies which delay or halt the progression of ESRD will provide a basis for the treatment of chronic renal disease. A variety of growth and differentiation factors, for example,

45 epidermal growth factor (EGF), transforming growth factor- α and - β (TGF- α and - β), insulin like growth factor-1 (IGF-1), fibroblast growth factor (FGF), platelet derived growth factor (PDGF) and bone morphogenetic protein (BMP) have been shown to participate in the regulation of the growth and repair of renal tissues. M.R. Hammerman and S.B. Miller, "Therapeutic Use of Growth Factors in Renal Failure," J. Am. Soc. Nephrol., 5: 1-11 (1994); and R.C. Harris, "Growth Factors and Cytokines in Acute Renal Failure," Adv. Renal. Repl. Ther., 4: 43-53 (1997).

50 **[0007]** Epidermal growth factor (EGF) enhances renal tubule cell regeneration and repair and accelerates the recovery of renal function in postischemic acute renal failure. H.D. Humes, D.A. Cieslinski, T. Coimbra, J.M. Messina and C. Galvao, J. Clin. Invest., 84: 1757-1761 (1989). Insulin-like growth factor I (IGF-1) accelerates recovery from ischemic acute tubular necrosis in the rat. S.B. Miller, D.R. Martin, J. Kissane, and M.R. Hammerman, Proc. Natl. Acad. Sci. USA, 89:11876-11880 (1994). Hepatocyte growth factor accelerates recovery from acute ischemic renal injury in rats. S.B. Miller, D.R. Martin, J. Kissane, and M.R. Hammerman, Am. J. Physiol., 266:129-134 (1994). Osteogenic protein-1 (bone morphogenetic protein-7 (BMP-7)) reduces severity of injury after ischemic acute renal failure in the rat. S. Vukicevic, V. Basic, D. Rogic, N. Basic, M.S. Shih, A. Shepard, D. Jin et al., J. Clin. Invest., 102: 202-214 (1998).

[0008] The term prostaglandin refers to compounds which are analogs of the natural prostaglandins PGD₁, PGD₂,

PGE₂, PGE₁ and PGF₂. These compounds bind to the prostaglandin receptors. Such binding is readily determined by those skilled in the art according to standard assays (e.g., S. An et al., Cloning and Expression of the EP₂ Subtype of Human Receptors for Prostaglandin E₂, Biochemical and Biophysical Research Communications, 197(1): 263-270 (1993)). These compounds can be synthesized by methods known in the art. See, e.g., Goodman and Gilman, The Pharmacological Basis of Therapeutics, Eighth Ed., Pergamon Press, pp. 601-604 (1990).

[0009] Prostaglandins are alicyclic compounds related to the basic compound prostanoic acid. The carbon atoms of the basic prostaglandin are numbered sequentially from the carboxylic carbon atom through the cyclopentyl ring to the terminal carbon atom on the adjacent side chain. Normally the adjacent side chains are in the trans orientation. The presence of an oxo group at C-9 of the cyclopentyl moiety is indicative of a prostaglandin within the E class while PGE₂ contains a trans unsaturated double bond at the C₁₃-C₁₄ and a cis double bond at the C₅-C₆ position. However, there are severe side effects associated with PGE₂ treatment. W.S.S. Jee and Y.F. Ma, Bone, 21:297-304 (1997).

[0010] A variety of prostaglandins are described and referenced below. However, other prostaglandins will be known to those skilled in the art.

[0011] U.S. Patent No. 4,177,346 discloses certain 1,5-disubstituted-2-pyrrolidones which are prostaglandin-like in chemical structure and have vasodilator activity, antihypertensive activity and antiseecretory activity.

[0012] International Patent Application, Publication No. WO 99/02164, discloses methods and compositions for treating impotence or erectile dysfunction by using prostaglandins that are selective EP₂ or EP₄ prostanoid receptor agonists.

[0013] U.S. Patent No. 4,112,236 discloses certain interphenylene 8-aza-9-dioxothia-11,12-secoprostaglandins which have renal vasodilatory activity and are useful for the treatment of patients with renal impairment. U.S. Patent No. 4,033,996 discloses certain 8-aza-9-oxo(and dioxo)-thia-11,12-secoprostaglandins which are useful as renal vasodilators, for the prevention of thrombus formation, to induce growth hormone release, and as regulators of the immune response.

[0014] Certain 11,12-secoprostaglandins and analogs thereof, which have a variety of therapeutic uses, including their use as renal vasodilators, are disclosed, for example, in the following: Great Britain Patent Nos. 1 478 281 and 1 479 156; U.S. Patent Nos. 3,987,091; 3,991,087; 3,991,106; 4,055,596; 4,066,692; and 4,175,203; J.B. Bicking et al., J. Med. Chem., 1977, Vol. 20, No. 1, pages 35-43; J.H. Jones et al., J. Med. Chem., 1977, Vol. 20, No. 1, pages 44-48; J.B. Bicking et al., J. Med. Chem., 1983, Vol. 26, pages 335-341; and J.B. Bicking et al., J. Med. Chem., 1983, Vol. 26, pages 342-348.

[0015] European Patent Application, Publication No. EP 0 911 321, and International Patent Applications, Publication Nos. WO 99/19300, WO 98/28264 and WO 98/ 58911, disclose certain prostaglandin agonists for the treatment of a variety of bone disorders, including osteoporosis, and for kidney degeneration.

[0016] Thus, there is a continuing need and a continuing search in this field of art for therapies for treating renal failure or dysfunction. More specifically, there is a need for receptor selective prostaglandin therapies, which do not have the side effects caused by non-selective agents.


SUMMARY OF THE INVENTION

[0017] The present invention particularly provides methods of treating acute or chronic renal failure or dysfunction, or a condition caused thereby, in a mammal comprising administering to said mammal an EP₄ receptor selective agonist, an isomer thereof, a prodrug of said agonist or isomer, or a pharmaceutically acceptable salt of said agonist, isomer or prodrug. More particularly, the present invention provides said methods wherein said condition is selected from the group consisting of hypertension, congestive heart failure, glomerulonephritis, uremia and chronic renal insufficiency.

[0018] Also, the present invention particularly provides methods of using a pharmaceutical composition comprising a compound which is a selective EP₄ receptor agonist, an isomer thereof, a prodrug of said agonist or isomer, or a pharmaceutically acceptable salt of said agonist, isomer or prodrug, together with a physiologically acceptable diluent, carrier or vehicle, for the treatment of acute or chronic renal failure or dysfunction, or a condition caused thereby, in a mammal. More particularly, the present invention provides said methods wherein said condition is selected from the group consisting of hypertension, congestive heart failure, glomerulonephritis, uremia and chronic renal insufficiency.


[0019] The term "prostaglandin agonist" refers to a compound, including its isomers, prodrugs and pharmaceutically acceptable salts, which binds to a prostaglandin receptor, such as EP₄, and mimics the action of the prostaglandin *in vivo*. A variety of such compounds are described and referenced above. However, other prostaglandin agonists will be known to those skilled in the art.

[0020] Preferred EP₄ selective agonists for use in the present invention include compounds of formula I:



prodrugs thereof or pharmaceutically acceptable salts of said compounds or said prodrugs, wherein:

- Q is COOR³, CONHR⁴ or tetrazol-5-yl;
 A is a single or cis double bond;
 B is a single or trans double bond;
 U is



- R² is α -thienyl, phenyl, phenoxy, monosubstituted phenyl or monosubstituted phenoxy, said substituents being selected from the group consisting of chloro, fluoro, phenyl, methoxy, trifluoromethyl and (C₁-C₃)alkyl;
 R³ is hydrogen, (C₁-C₅)alkyl, phenyl or p-biphenyl;
 R⁴ is COR⁵ or SO₂R⁵; and
 R⁵ is phenyl or (C₁-C₅)alkyl.

[0021] The compounds of formula I may be prepared as disclosed in commonly assigned U.S. Patent No. 4,177,346, which is hereby incorporated by reference herein.

[0022] A preferred group of EP₄ receptor selective agonists of formula I are those compounds of formula I wherein Q is 5-tetrazolyl. Particularly preferred compounds within this group include 5-(3-hydroxy-4-phenyl-but-1-enyl)-1-[6-(1H-tetrazol-5-yl)-hexyl]-pyrrolidin-2-one and 5-(3-hydroxy-4-phenyl-butyl)-1-[6-(1H-tetrazol-5-yl)-hexyl]-pyrrolidin-2-one.

[0023] Another preferred group of EP₄ receptor selective agonists of formula I are those compounds of formula I wherein Q is COOH. Particularly preferred compounds within this group include 7-[2-(3-hydroxy-4-phenyl-but-1-enyl)-5-oxo-pyrrolidin-1-yl]-heptanoic acid and 7-(2-(3-hydroxy-4-phenyl-butyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid.

[0024] The term "treating", "treat" or "treatment" as used herein includes preventative (e.g., prophylactic), palliative and curative treatment.

[0025] The term "pharmaceutically acceptable" means the carrier, vehicle, diluent, excipients, and/or salt must be compatible with the other ingredients of the formulation, and not deleterious to the recipient thereof.

[0026] The expression "prodrug" refers to a compounds that is a drug precursor which, following administration, releases the drug *in vivo* via some chemical or physiological process (e.g., a prodrug on reaching the physiological pH or through enzyme action is converted to the desired drug form). Exemplary prodrugs upon cleavage release the corresponding drug compounds.

[0027] The expression "pharmaceutically acceptable salt" refers to nontoxic anionic salts containing anions such as, but not limited to, chloride, bromide, iodide, sulfate, bisulfate, phosphate, acetate, maleate, fumarate, oxalate, lactate, tartrate, citrate, gluconate, methanesulfonate and 4-toluene-sulfonate. The expression also refers to nontoxic cationic salts such as, but not limited to, sodium, potassium, calcium, magnesium, ammonium or protonated benzathine (N,N'-dibenzylethylenediamine), choline, ethanolamine, diethanolamine, ethylenediamine, meglamine (N-methylglucamine), benethamine (N-benzylphenethylamine), piperazine and tromethamine (2-amino-2-hydroxymethyl-1,3-propanediol).

[0028] Other features and advantages will be apparent from the description and the appendant claims which describe the present invention.

DETAILED DESCRIPTION OF THE INVENTION

[0029] Any EP₄ receptor selective agonist may be used as the EP₄ receptor selective agonist of this invention. EP₄ selective agonists are compounds which have an IC₅₀ at the EP₁, EP₂ and EP₃ receptor which is at least 10-fold greater than the IC₅₀ at the EP₄ receptor subtype. For example, 7-(2-(3-hydroxy-4-phenyl-butyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid is an EP₄ receptor selective PGE₂ agonist with an EP₄ receptor binding IC₅₀ of 16 nM. At all other EP receptor subtypes, including the EP₁, EP₂ and EP₃ receptor subtypes, the IC₅₀ for 7-(2-(3-hydroxy-4-phenylbutyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid is greater than 3200 nM.

[0030] Accordingly, high selectivity or specificity for the EP₄ receptor, compared to other prostaglandin receptors, characterizes the compounds to be used in the methods and compositions according to the present invention. The more specific the compound is for the EP₄ receptor, the better are the therapeutic results that are obtained in the methods and compositions according to the present invention. Also, the receptor selectivity of the compounds to be used in the methods and compositions of the present invention results in the lessening or elimination of undesirable side effects caused by nonselective agents.

[0031] The EP₄ receptor selective agonists used in the methods of this invention can be adapted to therapeutic use in animals, e.g., mammals, and particularly humans.

These compounds exhibit renal vasodilatory activity, and, therefore, are useful for the treatment of patients with renal impairment. Included in this group are patients with hypertension, renal failure, diabetes-induced renal failure, congestive heart failure, glomerulonephritis, uremia, and chronic renal insufficiency. These compounds by virtue of their renal vasodilatory activity improve renal function.

[0032] The utility of the EP₄ selective agonists used in the methods of the present invention as medical agents in the treatment of acute or chronic renal failure or dysfunction in animals, e.g., mammals, especially humans, is demonstrated by the activity of those agonists in conventional assays, including the prostaglandin receptor binding assay, the cyclic AMP assay, and *in vivo* assays, including the acute renal failure assays, all of which are described below. Such assays also provide a means whereby the activities of the EP₄ selective agonists can be compared to each other and with the activities of other known compounds and compositions. The results of these comparisons are useful for determining dosage levels in animals, e.g., mammals, including humans, for the treatment of such diseases.

Determination of cAMP Elevation in 293-S Cell Lines Stably Overexpressing Recombinant Human EP₄ Receptors

[0033] cDNA representing the complete open reading frame of the human EP₄ receptor is generated by reverse transcriptase polymerase chain reaction using oligonucleotide primers based on published sequences (J.W. Regan, T.J. Bailey, D.J. Pepperl, K.L. Pierce, A.M. Bogardus, J.E. Donello, C.E. Fairbairn, K.M. Kedzie, D.F. Woodward and D.W. Gil, "Cloning of a Novel Human Prostaglandin Receptor with Characteristics of the Pharmacologically Defined EP₂ Subtype," *Mol. Pharmacology* 46:213-220 (1994)) and RNA from primary human lung cells (EP₄) as templates. cDNAs are cloned into the multiple cloning site of pcDNA3 (Invitrogen Corporation, 3985B Sorrento Valley Blvd., San Diego, CA 92121) and used to transfect 293-S human embryonic kidney cells via calcium phosphate coprecipitation. G418-resistant colonies are expanded and tested for specific [³H]PGE₂ binding. Transfectants demonstrating high levels of specific [³H]PGE₂ binding are further characterized by Scatchard analysis to determine B_{max} and K_ds for PGE₂. The lines selected for compound screening have approximately 256,400 receptors per cell and a K_d = 2.9 nM for PGE₂ (EP₄). Constitutive expression of the receptor in parental 293-S cells is negligible. Cells are maintained in RPMI supplemented with fetal bovine serum (10% final) and G418 (700 ug/ml final).

[0034] cAMP responses in the 293-S/EP₄ lines are determined by detaching cells from culture flasks in 1 ml of Ca⁺⁺ and Mg⁺⁺ deficient PBS via vigorous pounding, adding serum-free RPMI to a final concentration of 1 X 10⁶ cells/ml, and adding 3-isobutyl-1-methylxanthine (IBMX) to a final concentration of 1 mM. One milliliter of cell suspension is immediately aliquoted into individual 2 ml screwcap microcentrifuge tubes and incubated for 10 minutes, uncovered, at 37 °C, 5% CO₂, 95% relative humidity. The compound to be tested in DMSO or ethanol is then added to cells at 1:100 dilutions such that final DMSO or ethanol concentrations is 1%. Immediately after adding compound, the tubes are covered, mixed by inverting two times, and incubated at 37 °C for 12 minutes. Samples are then lysed by incubation at 100 °C for 10 minutes and immediately cooled on ice for 5 minutes. Cellular debris is pelleted by centrifugation at 1000 X g for 5 minutes, and cleared lysates are transferred to fresh tubes. cAMP concentrations are determined using a commercially available cAMP radioimmunoassay (RIA) kit (NEK-033, DuPont/NEN Research Products, 549 Albany St., Boston, MA 02118) after diluting cleared lysates 1:10 in cAMP RIA assay buffer (included in kit). Typically, the cells are treated with 6-8 concentrations of the compound to be tested in 1 log increments. EC₅₀ calculations are performed on a calculator using linear regression analysis on the linear portion of the dose response curves.

Assay for Binding to Prostaglandin E₂

[0035] Membrane Preparation: All operations are performed at 4 °C. Transfected cells expressing prostaglandin E₂ type 1 receptors (EP₁), type 2 (EP₂), type 3 (EP₃) or type 4 (EP₄) receptors are harvested and suspended to 2 million cells per ml in Buffer A [which is composed of the following: 50 mM Tris-HCl (pH 7.4), 10 mM MgCl₂, 1 mM EDTA, 1 mM Pefabloc peptide (Boehringer Mannheim Corp., Indianapolis, IN), 10 μM Phosphoramidon peptide (Sigma, St. Louis, MO), 1 μM pepstatin A peptide (Sigma, St. Louis, MO), 10 μM Melastatinal peptide (Sigma, St. Louis, MO), and 100 μM antipain peptide (Sigma, St. Louis, MO)]. The cells are lysed by sonification with a Branson Sonifier (Model #250, Branson Ultrasonics Corporation, Danbury, CT) in 2 fifteen second bursts. Unlysed cells and debris are removed by centrifugation at 100 x g for 10 min. Membranes are then harvested by centrifugation at 45,000 x g for 30 minutes. Pelleted membranes are resuspended to 3-10 mg of protein per ml of Buffer A, the protein concentration being determined by the method of Bradford [M. Bradford, Anal. Biochem., 72, 248 (1976)]. Resuspended membranes are then stored frozen at -80 °C until use.

[0036] Binding Assay: Frozen membranes prepared as above are thawed and diluted to 1 mg protein per ml in Buffer A above. One volume of membrane preparation is combined with 0.05 volume test compound or buffer and one volume of 3 nM ³H-prostaglandin E₂ (#TRK 431, Amersham, Arlington Heights, IL) in Buffer A. The mixture (205 μL total volume) is incubated for 1 hour at 25°C. The membranes are then recovered by filtration through type GF/C glass fiber filters (#1205-401, Wallac, Gaithersburg, MD) using a Tomtec harvester (Model Mach II/96, Tomtec, Orange, CT). The membranes with bound ³H-prostaglandin E₂ are trapped by the filter, while the buffer and unbound ³H-prostaglandin E₂ pass through the filter into waste. Each sample is then washed 3 times with 3 ml of 50 mM Tris-HCl (pH 7.4), 10 mM MgCl₂, and 1 mM EDTA. The filters are then dried by heating in a microwave oven. To determine the amount of ³H-prostaglandin bound to the membranes, the dried filters are placed into plastic bags with scintillation fluid and counted in a LKB 1205 Betaplate reader (Wallac, Gaithersburg, MD). IC₅₀s are determined from the concentration of test compound required to displace 50% of the specifically bound ³H-prostaglandin E₂.

[0037] The full length coding sequence for the EP₁ receptor is made as disclosed in Funk et al., Journal of Biological Chemistry, 1993, 268, 26767-26772. The full length coding sequence for the EP₂ receptor is made as disclosed in Regan et al., Molecular Pharmacology, 1994, 46, 213-220. The full length coding sequence for the EP₃ receptor is made as disclosed in Regan et al., British Journal of Pharmacology, 1994, 112, 377-385. The full length coding sequence for the EP₄ receptor is made as disclosed in Bastien, Journal of Biological Chemistry, 269: 11873-11877 (1994). These full length receptors are used to prepare 293S cells expressing the EP₁, EP₂, EP₃ and EP₄ receptors. Additional information on the human EP₂ and human EP₄ prostaglandin receptors is disclosed in U.S. Patent Nos. 5,605,814; 5,716,835; and 5,759,789; which are hereby incorporated by reference herein.

[0038] 293S cells expressing one of the human EP₁, EP₂, EP₃ or EP₄ prostaglandin E₂ receptors are generated according to methods known to those skilled in the art. Typically, PCR (polymerase chain reaction) primers corresponding to the 5' and 3' ends of the published full length receptor are made according to the well known methods disclosed above and are used in an RT-PCR reaction using the total RNA from human kidney (for EP₁), human lung (for EP₂), human lung (for EP₃) or human lymphocytes (for EP₄) as a source. PCR products are cloned by the TA overhang method into pCR2.1 (Invitrogen, Carlsbad, CA) and identity of the cloned receptor is confirmed by DNA sequencing.

[0039] 293S cells (Mayo, Dept. of Biochemistry, Northwestern Univ.) are transfected with the cloned receptor in pcDNA3 (Invitrogen Corporation, 3985B Sorrento Valley Blvd., San Diego, CA 92121) by electroporation. Stable cell lines expressing the receptor are established following selection of transfected cells with G418.

[0040] Clonal cell lines expressing the maximal number of receptors are chosen following a whole cell ³H-PGE₂ binding assay using unlabeled PGE₂ as a competitor.

Acute Renal Failure Rat Model

[0041] The remnant rat kidney (5/6 nephrectomy) model employed was essentially as previously described in the literature (Vukicevic et al., Journal of Bone and Mineral Research, 2:533-545 (1987)). In brief, male rats (4 months old, weighing approximately 400 g) were anesthetized by intraperitoneal administration of ketamin (20 mg/kg) and then subjected to unilateral nephrectomy (left kidney). All animals were subjected to intraperitoneal administration of 1-3 ml of prewarmed saline to compensate for any fluid loss during the surgery. After one week, 2/3 of the remaining kidney was surgically removed. The rats were allowed to recover for one or two weeks prior to the initiation of the therapy for the prevention of the disease. The rats were pair fed to comparable weight for control and treated animals. The 5/6 nephrectomy model has been shown to reproduce several features of progressive human renal disease; namely, glomerulosclerosis, tubulointerstitial disease, proteinuria, progressive decline in kidney function and the development of systemic hypertension.

[0042] Blood samples (0.5 ml) were obtained from the orbital plexus and serum creatinine, BUN (blood urea nitrogen), calcium, phosphate and other blood chemistries were monitored throughout the study. Urine was collected on metabolic

cages for 24 h. Creatinine was measured by a Jaffe method. A.W. Whelton, A.J. Watson and R.C. Rock: "Nitrogen Metabolites and Renal Function," in: Clinical Chemistry (Eds: C.A. Burtis and E.R. Ashwood) W.B. Saunders, Philadelphia, PA, 1513-1575 (1994). BUN was measured by a glutamate dehydrogenase ultraviolet method, phosphorus by a molybdate method, and calcium by an o-cresolphthaleine method. R.A. Lafayette, R.D. Perrone and A.S. Levey: "Laboratory Evaluation of Renal Function," in: Diseases of the Kidney (Eds: R.W. Schrier and C.W. Gottschalk), Little, Brown and Company, Inc., Vol. 6, 307-354 (1997). The GFR was calculated using serum creatinine over urine creatinine (24 hr collection) and adjusted to body weights. The cumulative survival rate was also observed and recorded for both the control and the experimental groups. The kidney was subjected to histomorphometric analysis and scored for tissue pathology.

[0043] Results: The efficacy of 7-(2-(3-hydroxy-4-phenyl-butyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid, an EP₄ receptor selective PGE₂ agonist of the present invention, was tested in pre-clinical models of acute renal failure. Two different models of acute renal failure were used: the nephrotoxic acute renal failure model, as described above, and the reperfusion renal injury model, which involved clamping both renal arteries for 60 minutes. (K.J. Kelly, W.W. Williams, R.B. Colvin and J.V. Bonventure, "Intercellular Adhesion Molecule 1 Protects the Kidney Against Ischemic Injury," Proc. Natl. Acad. Sci. USA, 91:812-816 (1994); and S. Vukicevic, V. Basic, D. Rogic, N. Basic, M.S. Shih, A. Shepard, D. Jin, D. Bosukonda, W. Jones, H. Dorai, S. Ryan, D. Griffiths, J. Maliakal, M. Jelic, M. Pastorcic, A. Stavljenic and K. Sampath, "Osteogenic Protein-1 (Bone Morphogenetic Protein-7) Reduces Severity of Injury After Ischemic Acute Renal Failure in Rat," J. Clin. Invest., 102:202-214 (1998).)

[0044] In the nephrotoxic acute renal failure model, the EP₄ agonist was effective in protecting against renal injury, based on serum parameters, and increased the survival time of animals in a dose dependent manner, with a dose of 10 mg/kg being better than the dose of 1 mg/kg. In this study, four out of ten animals survived in the control group after seven days, whereas nine out of ten animals survived in the treated group at a dose of 10 mg/kg. In the reperfusion renal injury model, the EP₄ agonist was effective 24 hours following reperfusion. The animals were dosed with the EP₄ agonist just prior to the initiation of the nephrotoxic insult.

[0045] Administration of an EP₄ receptor selective agonist according to the methods of this invention can be via any available mode which delivers the EP₄ receptor selective agonist systemically or locally. These methods include oral routes, parenteral, intraduodenal routes, etc. Generally, the compounds of this invention are administered orally, but parenteral administration (e.g., intravenous, intramuscular, transdermal, subcutaneous, rectal or intramedullary) may be utilized, for example, where oral administration is inappropriate for the target or where the patient is unable to ingest the drug.

[0046] In any event, the amount and timing of the compound administered will, of course, be dependent on the subject being treated, on the severity of the affliction, on the manner of administration and on the judgment of the prescribing physician. Thus, because of patient to patient variability, the dosages given herein are a guideline and the physician may titrate doses of the drug compound to achieve the treatment that the physician considers appropriate for the patient. In considering the degree of treatment desired, the physician must balance a variety of factors such as age of the patient, presence of preexisting disease, as well as presence of other diseases (e.g., cardiovascular disease).

[0047] The EP₄ receptor selective agonist compounds used in the methods of this invention are generally administered in the form of a pharmaceutical composition comprising at least one of the compounds of this invention together with a pharmaceutically acceptable vehicle or diluent. Thus, the EP₄ receptor selective agonist compound can be administered individually in any conventional form, such as oral, intranasal, parenteral, rectal or transdermal dosage form.

[0048] For oral administration the pharmaceutical composition can take the form of solutions, suspensions, tablets, pills, capsules, powders, and the like. Tablets containing various excipients such as sodium citrate, calcium carbonate and calcium phosphate are employed along with various disintegrants such as starch, preferably potato or tapioca starch, and certain complex silicates, together with binding agents such as polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type are also employed as fillers in soft and hard-filled gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the compositions of this invention can be combined with various sweetening agents, flavoring agents, coloring agents, emulsifying agents and/or suspending agents, as well as such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

[0049] The compounds can also be administered orally in solid solution with lipids such as cholesterol acetate. The inclusion of lipid in the formulation markedly increases absorption of the compound or analog. Preparation of such formulations is described in detail in Rudel, U.S. Patent No. 3,828,106, which is incorporated herein by reference.

[0050] For purposes of parenteral administration, solutions in sesame or peanut oil or in aqueous propylene glycol can be employed, as well as sterile aqueous solutions of the corresponding water-soluble salts. Such aqueous solutions may be suitably buffered, if necessary, and the liquid diluent first rendered isotonic with sufficient saline or glucose.

These aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal injection purposes. In this connection, the sterile aqueous media employed are all readily obtainable by standard techniques well-known to those skilled in the art.

[0051] Compositions to be administered intravenously or by injection can be prepared as solutions of the compound in, for example, an isotonic aqueous solution, an alcohol solution, an ethanol-saline solution, or an ethanol-dextrose solution. Ethanol can be added to the solution to increase solubility and other additives such as methylparaben or other ingredients such as fillers, colorings, flavorings, diluents and the like can be included. The composition can also be administered as a suspension of the compound or analog in aqueous or nonaqueous media.

[0052] Among the preferred formulations for administration intravenously or by injection are complexes of the active ingredient with α -cyclodextrin. Preparation of complexes of compounds and analogs with α -cyclodextrin clathrates are described in detail in Hayashi et al., U.S. Patent No. 4,054,736, which is incorporated herein by reference. Complexes wherein the ratio of α -cyclodextrin to a compound of this invention is 97:3 are especially preferred.

[0053] For purposes of transdermal (e.g., topical) administration, dilute sterile, aqueous or partially aqueous solutions (usually in about 0.1% to 5% concentration), otherwise similar to the above parenteral solutions, are prepared.


[0054] Methods of preparing various pharmaceutical compositions with a certain amount of active ingredient are known, or will be apparent in light of this disclosure, to those skilled in the art. For examples of methods of preparing pharmaceutical compositions, see Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., 19th Edition (1995). Thus, as described above, while the compounds of this invention may be administered to the patients in any of the known formulations or modes of administration.

[0055] A preferred dosage for the methods and compositions of the present invention is about 0.001 to about 100 mg/kg/day of an EP₄ receptor selective agonist, an isomer thereof, a prodrug of said agonist or isomer, or a pharmaceutically acceptable salt of said agonist, isomer or prodrug. An especially preferred dosage is about 0.01 to about 10 mg/kg/day of an EP₄ receptor selective agonist, an isomer thereof, a prodrug of said agonist or isomer or a pharmaceutically acceptable salt of said agonist, isomer or prodrug.

[0056] More specifically, for administration intravenously or by injection, the dosages of the compounds to be used in the methods and compositions of the present invention can range from about 5 to about 60 ng/min/kg body weight, with the preferred dosage range being from about 10 ng/kg/min to about 30 ng/kg/min. If the dose is administered by intravenous injection; it should not exceed about 100 μ g/kg body weight per day.

Claims

1. Use of an EP₄ receptor selective agonist, an isomer thereof, a prodrug of said agonist or isomer, or a pharmaceutically acceptable salt of said agonist, isomer or prodrug, for the manufacture of a medicament for treating acute or chronic renal failure or dysfunction, or a condition caused thereby, in a mammal.
2. The use of claim 1 wherein said condition is selected from the group consisting of hypertension, congestive heart failure, glomerulonephritis, uremia and chronic renal insufficiency.
3. The use of claim 1 wherein said EP₄ selective agonist is a compound of formula I:



a prodrug thereof or a pharmaceutically acceptable salt of said compound or said prodrug, wherein:

Q is COOR³, CONHR⁴ or tetrazol-5-yl;

A is a single or cis double bond;

B is a single or trans double bond;
U is



10 R^2 is α -thienyl, phenyl, phenoxy, monosubstituted phenyl or monosubstituted phenoxy, said substituents being selected from the group consisting of chloro, fluoro, phenyl, methoxy, trifluoromethyl and ($\text{C}_1\text{-C}_3$)alkyl;

R^3 is hydrogen, ($\text{C}_1\text{-C}_5$)alkyl, phenyl or p-biphenyl;

R^4 is COR^5 or SO_2R^5 ; and

R^5 is phenyl or ($\text{C}_1\text{-C}_5$)alkyl.

- 15 4. The use of claim 3 wherein said EP_4 receptor selective agonist is a compound of formula 1 wherein Q is 5-tetrazolyl.
5. The use of claim 4 wherein said compound is 5-(3-hydroxy-4-phenyl-but-1-enyl)-1-[6-(1 H-tetrazol-5-yl)-hexyl]-pyrrolidin-2-one or 5-(3-hydroxy-4-phenyl-butyl)-1-[6-(1 H-tetrazol-5-yl)-hexyl]-pyrrolidin-2-one.
- 20 6. The use of claim 3 wherein said EP_4 receptor selective agonist is a compound of formula I wherein Q is COOH .
7. The use of claim 6 wherein said compound is 7-(2-(3-hydroxy-4-phenyl-butyl)-5-oxo-pyrrolidin-1-yl)-heptanoic acid or 7-[2-(3-hydroxy-4-phenyl-but-1-enyl)-5-oxo-pyrrolidin-1-yl]-heptanoic acid.
- 25 8. A pharmaceutical composition for the treatment of acute or chronic renal failure or dysfunction, or a condition caused thereby, in a mammal comprising a compound which is a selective EP_4 receptor agonist, an isomer thereof, a prodrug of said agonist or isomer, or a pharmaceutically acceptable salt of said agonist, isomer or prodrug, and a pharmaceutically acceptable diluent, carrier or vehicle.
- 30 9. The composition of claim 8 wherein said condition is selected from the group consisting of hypertension, congestive heart failure, glomerulonephritis, uremia and chronic renal insufficiency.
10. The composition of claim 8 wherein said EP_4 selective agonist is a compound defined in any one of claims 3 to 7.